

TCN News

News and announcements from the Tobacco Control Network

TCN Announcements

 A new <u>narrated presentation</u> from TCN Chair Frances Limtiaco provides an overview of the 2019-2022 TCN Strategic Map. The video presentation is also available on the <u>Resources tab</u> of the TCN website.

Happy Anniversary!

In the first newsletter of each month, TCN recognizes anniversaries of state and territorial tobacco program managers to thank them for their leadership in tobacco control and public health more broadly. Thank you and congratulations to the following program managers!

- Jennifer Ullman (MT) 11 years.
- Elizabeth Guerrero (GU) 9 years.

Partner Announcements

 FDA announced that they filed a federal court motion to extend the deadline for tobacco product premarket review applications by 120 days from May 12 to Sept. 9, citing the COVID-19 outbreak. The twice monthly **TCN News** provides a concise
digest of tobacco control
announcements to the
Tobacco Control Network's
membership base of tobacco
control program managers and
additional staff from each
state, territory, and D.C., as
well as our valued partners.

Upcoming Events

Research into Action: Why
Point of Sale Matters —
Community Anti-Drug
Coalitions of America
Webinar
April 16, 1-2 p.m. ET
| Registration required

Request for Information on Vaping Products Associated with Lung Injuries – FDA April 20 | Comment deadline

Increasing Lung Cancer
Screening Uptake in Eligible
Adults: Do We Know What
Works? – American Cancer
Society
Webinar
April 29, 1-2 p.m. ET
| Registration required

No Menthol Sunday – National African American Tobacco Prevention Network May 17 | Save the date

Tobacco Cessation and Lung

- CDC researchers published the following articles:
 - A Journal of Smoking Cessation analysis of national quitline data from 2012 to 2015 found that the average call lasted 11.4 minutes, and that there was no relationship between quitline service provider and the volume or duration of calls in a given state.
 - o In a Preventing Chronic Disease analysis of SummerStyles survey data, 62.6% of adults agreed that nicotine in e-cigarette products is harmful to the adolescent brain, with significant variance observed across different subpopulations.
- Articles from <u>Truth Initiative</u> and the <u>American Nonsmokers' Rights Foundation</u> explore the possible link between tobacco use and poorer outcomes of COVID-19 infection.
- Public Health Law Center <u>updated</u> their listing of state laws that prohibit smoking in cars with children present.
- The Duke-UNC Tobacco Treatment Specialist
 Training Program announced that their program will now be available virtually through a combination of video classes and self-paced online trainings.

State News

- The Minnesota Department of Health <u>posted</u> a summary of findings related to smoking, vaping, lung health, and COVID-19.
- McCarran Airport in Las
 Vegas, Nevada announced that they closed their
 indoor smoking lounges.
- According to a <u>poll</u> commissioned by the Campaign for Tobacco-Free Kids, 70% of **New**

Cancer Screening – American Cancer Society Webinar June 3, 1-2 p.m. ET | Registration required

State of Black Health
Conference – National African
American Tobacco Prevention
Network
Charlotte, NC
Sept. 1-3 | Registration
required

Nicotine Dependence
Conference & National Summit
on Smokeless Tobacco
Prevention 2020 – Mayo Clinic
Rochester, MN
Oct. 18-20 | Registration
required

Funding Opportunities

If you are aware of any funding opportunities that would appeal to state and territorial tobacco control programs, please reach out to tcn@astho.org.

Job Listings

Senior Science Advisor FDA Center for Tobacco Products Silver Spring, MD

Health Communications
Specialist
California Department of
Public Health
Sacramento, CA

Policy Consultant
California Department of
Public Health
Sacramento, CA

About the Tobacco Control Network

The TCN's mission is to improve the public's

York residents <u>support</u> a policy that would end the sale of flavored e-cigarette products.

A Tobacco Control microsimulation model
 of Minnesota from 2018 to 2037 estimates that
 over 20 years, projected smoking prevalence
 decreases will result in 14,063 fewer smoking related deaths and nearly \$20 billion in reduced
 healthcare spending.

In the News

- According to documents uncovered by AP, JUUL Labs <u>lobbied</u> to prevent state attorneys general from filing lawsuits against the company, and enlisted lowa Attorney General Tom Miller as an advisor.
- NYC CLASH, a smokers rights group, lost a federal lawsuit against the U.S. Department of Housing and Urban Development in an attempt to overturn the agency's smoke-free housing policy.

Research

- A Tobacco Control analysis of 2014-2017 National Youth Tobacco Survey data <u>found</u> that less than 1% of U.S. adolescents who use e-cigarettes were previously established cigarette smokers.
- An online experiment of 16 FDA-developed pictorial cigarette warnings published in *Nicotine & Tobacco Research* concluded that a majority of the warnings resulted in greater consideration of smoking risks compared to a control of current Surgeon General's warnings.

health by providing education and statebased expertise to tobacco prevention and control at the state and national levels.

The TCN is a network of tobacco program managers and staff from state and territorial health departments as well as Funder's Alliance organizations, and a peer group of the Association of State and Territorial Health Officials (ASTHO) on behalf of the CDC's Office on Smoking and Health.

About ASTHO

The Association of State and Territorial Health Officials is the national nonprofit association representing the state and territorial public health agencies of the United States, the U.S. Territories, and the District of Columbia. ASTHO members, the chief health officials of these jurisdictions, are dedicated to formulating and influencing sound public health policy and to ensuring excellence in state-based public health practice.



 A Nicotine & Tobacco Research analysis of college student tobacco use data <u>concluded</u> that students fell into four tobacco use trajectories abstainers/dabblers, adult smokers, college smokers, and teenage users—and identified risk factors for each of those trajectories.

Feel free to forward this communication or encourage others to subscribe directly.

This message was intended for << Test Email Address >>. If you believe this has been sent to you in error, please <u>unsubscribe</u> or <u>update your subscription preferences</u>.

Our address:

The Association of State and Territorial Health Officials, 2231 Crystal Drive, Suite 450, Arlington, VA 22202 TCN@astho.org

